

The Arthur Terry Learning Partnership

Kittoe Road, Sutton Coldfield West Midlands, B74 4RZ Tel: (0121) 323 1134 Fax: (0121) 308 8033

...

Richard Gill CBE NPQH, NLE, FRSA Chief Executive Officer (CEO)

Dear Parent / Carer,

## Re: Increase in scarlet fever

We are writing to you as you may be aware of reports of the national increase in notifications of scarlet fever to the UK Health Security Agency (UKHSA), above seasonal expected levels.

Although not all areas in the UK are experiencing the same levels of cases, we would like to take this opportunity to remind you of the signs, symptoms and the actions to be taken if you think that you or your child might have scarlet fever, by sharing the following UKHSA information with you.

## Signs and symptoms of scarlet fever

Scarlet fever is a common childhood infection caused by *Streptococcus pyogenes*, or group A Streptococcus (GAS). It is not usually serious, but should be treated with antibiotics to reduce the risk of complications (such as pneumonia) and spread to others. The early symptoms of scarlet fever include sore throat, headache, fever, nausea and vomiting. After 12 to 48 hours, the characteristic red, pinhead rash develops, typically first appearing on the chest and stomach, then rapidly spreading to other parts of the body, and giving the skin a sandpaper-like texture. The scarlet rash may be harder to spot on darker skin, although the 'sandpaper' feel should be present. Patients typically have flushed cheeks and be pale around the mouth. This may be accompanied by a bright red red 'strawberry' tongue.

If you think you, or your child, might have scarlet fever:

- contact your GP or NHS 111 as soon as possible
- make sure that you or your child take(s) the full course of any antibiotics prescribed. Although you
  or your child will feel better soon after starting the course of antibiotics, you must complete the
  course to ensure that you do not carry the bacteria in your throat after you have recovered
- stay at home, away from nursery, school or work for at least 24 hours after starting the antibiotic treatment, to avoid spreading the infection

You can help stop the spread of infection through frequent hand washing and by not sharing eating utensils, clothes, bedding and towels. All contaminated tissues should be disposed of immediately.

## Invasive Group A Strep (iGAS)

The same bacteria which cause scarlet fever can also cause a range of other types of infection such as skin infections (impetigo) and sore throat. In very rare cases, the bacteria can get into the bloodstream and cause an illness called invasive group A strep (iGAS). Whilst still very uncommon, there has been an increase in iGAS cases this year, particularly in children under 10 years old. It is very rare for children with scarlet fever to develop iGAS infection.

The Arthur Terry Learning Partnership, a charitable company limited by guarantee, registered in England and Wales, company number 07730920. Registered office: The Arthur Terry School, Kittoe Road, Four Oaks, Sutton Coldfield, West Midlands, B74 4RZ

As a parent, you should trust your own judgement.

Contact NHS 111 or your GP if:

- your child is getting worse
- your child is feeding or eating much less than normal
- your child has had a dry nappy for 12 hours or more or shows other signs of dehydration
- your baby is under 3 months and has a temperature of 38C, or is older than 3 months and has a temperature of 39C or higher
- your baby feels hotter than usual when you touch their back or chest, or feels sweaty
- your child is very tired or irritable

Call 999 or go to A&E if:

- your child is having difficulty breathing you may notice grunting noises or their tummy sucking under their ribs
- there are pauses when your child breathes
- your child's skin, tongue or lips are blue
- your child is floppy and will not wake up or stay awake

## Stop the spread

During periods of high incidence of scarlet fever, there may also be an increase in outbreaks in schools, nurseries and other childcare settings. Children with suspected scarlet fever should stay off nursery / school until **24 hours** after the start of appropriate antibiotic treatment.

Good hygiene practice such as hand washing remains the most important step in preventing and controlling spread of infection.

You may find the following NHS and Government resources helpful.

<u>NHS – Scarlet Fever</u> <u>Scarlet fever: symptoms, diagnosis and treatment</u>

Yours faithfully

Mrs S Kibble Headteacher Arthur Terry School